

2014

Herbivory in the Northern Curly-tailed Lizard (*Leiocephalus carinatus*)

Bonnie K. Kircher

Trinity University, kircherbk@gmail.com

Christopher D. Robinson

Trinity University, crobins3@trinity.edu

Michele A. Johnson

Trinity University, mjohnso9@trinity.edu

Follow this and additional works at: http://digitalcommons.trinity.edu/bio_faculty

Repository Citation

Kircher B. K., Robinson C. D., & Johnson M. A. (2014). Herbivory in the Northern Curly-tailed Lizard (*Leiocephalus carinatus*). *Caribbean Herpetology* 50:1–2.

This Article is brought to you for free and open access by the Biology Department at Digital Commons @ Trinity. It has been accepted for inclusion in Biology Faculty Research by an authorized administrator of Digital Commons @ Trinity. For more information, please contact jcostanz@trinity.edu.

Herbivory in the Northern Curly-tailed Lizard (*Leiocephalus carinatus*)

Bonnie K. Kircher*, Christopher D. Robinson, and Michele A. Johnson

Trinity University, Department of Biology, San Antonio, Texas 78212, USA.

*Corresponding author (kircherbk@gmail.com)

Date of publication: 19 May 2014.

Citation: Kircher BK, Robinson CD, & Johnson MA. (2014) Herbivory in the Northern Curly-tailed Lizard (*Leiocephalus carinatus*). *Caribbean Herpetology* 50:1–2.

Leiocephalus carinatus (Northern curly-tailed lizard) is distributed throughout the Cayman Islands, Cuba, and the Bahamas, and has been introduced in southern Florida (Schwartz & Henderson 1991). Most species of *Leiocephalus*, and *L. carinatus* in particular, have an opportunistic diet (Henderson & Powell 2009). One important component of the curly-tailed lizard diet is arthropods—primarily ants, beetles, cockroaches, and lepidopterans (Schoener *et al.* 1982; reviewed in Henderson & Powell 2009). Curly-tailed lizards are also known predators of smaller lizards, particularly anoles (e.g., Schoener *et al.* 2002). In addition, their diet is comprised of up to 47% plant matter, most commonly fruits and flowers (Schoener *et al.* 1982), and Grant (1940) reported one observation of *L. carinatus* consuming flowers of the Beach Morning Glory (*Ipomoea pescaprae*) in the Cayman Islands. Despite the unusually high concentration of plant matter in the diet of these lizards, documented observations of herbivory in this group remain relatively rare.

On 21–23 July 2013 we observed two separate events of herbivory in *L. carinatus*, in which individual adults were eating flowers from the Beach Morning Glory (*Ipomoea pescaprae*) on a private beach on Crooked Island, Bahamas (22.773 N, -74.205 W). Each lizard picked off parts of the flower petals and masticated and swallowed them, feeding for about five minutes each. On 23 July 2013, a third adult *L. carinatus* was observed in the same location with a wilted morning glory flower in its mouth (Fig. 1).

Herbivory in lizards most commonly evolves in species that occur on small islands, and is thought to result from a lower abundance of available arthropod prey in those habitats (Schoener *et al.* 1982; Cooper & Vitt 2002; Dutra *et al.* 2011). Further, Schoener *et al.* (1982) and Cooper & Vitt (2002) found that larger lizards are more likely to consume plant matter than smaller lizards, perhaps because of the greater caloric needs associated with larger body size. Our observation of flower consumption by large *L. carinatus* on the small (148 km²) Crooked Island is consistent with both of these previously described patterns.



Figure 1. *Leiocephalus carinatus* (A) feeding on flowers (B) of the Beach Morning Glory (*Ipomoea pescaprae*).

References

- Cooper WE, & Vitt LJ. (2002) Distribution, extent, and evolution of plant consumption by lizards. *Journal of Zoology*, 257, 487-517.
- Dutra GF, Siqueira CC, Vrcibradic D, Kiefer MC, & Rocha CFD. (2011) Plant consumption of insular and mainland populations of a tropical lizard. *Herpetologica*, 67, 32-45.
- Grant, C. (1940) The herpetology of the Cayman Islands. *Bulletin of the Institute of Jamaica*, 2, 1-65.
- Henderson RW, & Powell R. (2009) *Leicophalus carinatus* Gray, 1827, pp. 138-140. In: *Natural History of West Indian Amphibians and Reptiles*. University Press of Florida, Gainesville.
- Schoener TW, Slade JB, & Stinson CH. (1982) Diet and sexual dimorphism in the very catholic lizard genus, *Leiocephalus* of the Bahamas. *Oecologia*, 53, 160-169.
- Schoener TW, Spiller DA, & Losos JB. (2002) Predation on a common *Anolis* lizard: can the food-web effects of a devastating predator be reversed? *Ecological Monographs*, 72, 383-407.
- Schwartz A, & Henderson RW. (1991) *Amphibians and Reptiles of the West Indies: Description, Distributions, and Natural History*. University of Florida Press, Gainesville, 423 pp.